

Computads for Generalised Signatures

Ioannis Markakis ¹

YaMCATS 30



UNIVERSITY OF
CAMBRIDGE

**ONASSIS
FOUNDATION**

¹The speaker is being partially supported by the Onassis Foundation
Scholarship ID: F ZQ 039-1/2020-2021

Table of Contents

1 Motivation

2 Definitions

3 Examples

4 Results

Question

What does it mean for a mathematical structure to be free?

Question


What does it mean for a mathematical structure to be free?

$$\mathcal{C} \begin{array}{c} \xrightarrow{F_M} \\ \xleftarrow{U_M} \end{array} \text{Alg}_M$$

- Structures are M -algebras and free are in the image of F_M .

Question


What does it mean for a mathematical structure to be free?



- Structures are M -algebras and free are in the image of F_M .
- For some M , there are more general generating data for free algebras.

Question

What does it mean for a mathematical structure to be free?




- Structures are M -algebras and free are in the image of F_M .
- For some M , there are more general generating data for free algebras.
 - Free strict 2-category monad²

²Street, "Limits indexed by category-valued 2-functors"

Question

What does it mean for a mathematical structure to be free?




- Structures are M -algebras and free are in the image of F_M .
- For some M , there are more general generating data for free algebras.
 - Free strict 2-category monad²
 - Free ω -category monad³

²Street, “Limits indexed by category-valued 2-functors”

³Batanin, “Computads for finitary monads on globular sets”

Question

What does it mean for a mathematical structure to be free?




- Structures are M -algebras and free are in the image of F_M .
- For some M , there are more general generating data for free algebras.
 - Free strict 2-category monad²
 - Free ω -category monad³
 - Monads on globular sets³

²Street, “Limits indexed by category-valued 2-functors”

³Batanin, “Computads for finitary monads on globular sets”

Question

What does it mean for a mathematical structure to be free?



- Structures are M -algebras and free are in the image of F_M .
- For some M , there are more general generating data for free algebras.
 - Free strict 2-category monad²
 - Free ω -category monad³
 - Monads on globular sets³
 - **Monads over other presheaf topoi**

²Street, “Limits indexed by category-valued 2-functors”

³Batanin, “Computads for finitary monads on globular sets”


Computads for 2-categories



- A **2-graph** X is a diagram satisfying the *globularity conditions*

$$ss = st$$

$$ts = tt$$



Computads for 2-categories

$$\begin{array}{ccc} X_2 & \hookrightarrow & X_2^* \\ \begin{array}{c} \downarrow s \\ \downarrow t \\ \downarrow \\ \downarrow \end{array} & & \begin{array}{c} \downarrow s \\ \downarrow t \\ \downarrow \\ \downarrow \end{array} \\ X_1 & \hookrightarrow & X_1^* \\ \begin{array}{c} \downarrow s \\ \downarrow t \\ \downarrow \\ \downarrow \end{array} & & \begin{array}{c} \downarrow s \\ \downarrow t \\ \downarrow \\ \downarrow \end{array} \\ X_0 & \equiv & X_0 \end{array}$$


- A 2-graph X is a diagram satisfying the *globularity conditions*

$$ss = st \qquad ts = tt$$

- The free **2-category** on X consists of formal composite and coherence cells quotiented by the laws of 2-categories.


- A 2-graph X is a diagram satisfying the *globularity conditions*

$$ss = st \quad ts = tt$$



- The free 2-category on X consists of formal composite and coherence cells quotiented by the laws of 2-categories.

- A 2-graph X is a diagram satisfying the *globularity conditions*




$$ss = st$$

$$ts = tt$$

- The free 2-category on X consists of formal composite and coherence cells quotiented by the laws of 2-categories.

Computads for 2-categories




- A **2-computad** X is a diagram satisfying the *globularity conditions*

$$ss = st \quad ts = tt$$

where X_1^* is the set of formal composites from X_1 .

- The free 2-category on X consists of formal composite and coherence cells quotiented by the laws of 2-categories.

Computads for 2-categories



- A 2-computad X is a diagram satisfying the *globularity conditions*

$$ss = st \quad ts = tt$$

where X_1^* is the set of formal composites from X_1 .

- The free **2-category** on X consists of formal composite and coherence cells quotiented by the laws of 2-categories.

Question

Monads are mathematical structures. When is a monad free?

Question

Monads are mathematical structures. When is a monad free?

Idea

It should correspond to a theory with no equations when $\mathcal{C} = [\mathcal{I}^{\text{op}}, \text{Set}]$.

- weak ω -categories
- algebraic Kan complexes / quasicategories

Question

Monads are mathematical structures. When is a monad free?

Idea

It should correspond to a theory with no equations when $\mathcal{C} = [\mathcal{I}^{\text{op}}, \text{Set}]$.

- weak ω -categories
- algebraic Kan complexes / quasicategories

Definition

A signature⁴ over $\mathcal{C} = [\mathcal{I}^{\text{op}}, \text{Set}]$ is a presheaf $\Sigma \in \mathcal{C}$ of function symbols with arity functions $B_{\bullet} : \Sigma_i \rightarrow \text{ob } \mathcal{C}$ satisfying for $\delta : j \rightarrow i$ that

$$B_{\delta * f} = B_f$$

⁴Bourke and Garner, “Monads and theories”

Definition

A Σ -algebra \mathbb{X} is a presheaf X equipped with functions

$$f^{\mathbb{X}} : \mathcal{C}(B_f, X) \rightarrow X_i$$

for $f \in \Sigma_j$ satisfying that $\delta^* \circ f^{\mathbb{X}} = (\delta^* f)^{\mathbb{X}}$ for $\delta : j \rightarrow i$.

- The forgetful functor $U_{\Sigma} : \text{Alg}_{\Sigma} \rightarrow \mathcal{C}$ is strict monadic.
- The free monad on Σ is $M_{\Sigma} = U_{\Sigma} F_{\Sigma}$.

Definition

A Σ -algebra \mathbb{X} is a presheaf X equipped with functions

$$f^{\mathbb{X}} : \mathcal{C}(B_f, X) \rightarrow X_i$$

for $f \in \Sigma_i$ satisfying that $\delta^* \circ f^{\mathbb{X}} = (\delta^* f)^{\mathbb{X}}$ for $\delta : j \rightarrow i$.

- The forgetful functor $U_{\Sigma} : \text{Alg}_{\Sigma} \rightarrow \mathcal{C}$ is strict monadic.
- The free monad on Σ is $M_{\Sigma} = U_{\Sigma} F_{\Sigma}$.

Problem

Weak ω -categories are not algebras of a signature, since the source of the associator is not a 1-dimensional function symbol, but a composite of them.

Table of Contents

1 Motivation

2 Definitions

3 Examples

4 Results

Definition

A direct category \mathcal{I} is a small category equipped with a dimension function $\dim : \text{ob } \mathcal{I} \rightarrow \text{Ord}$ to the class of ordinals such that $\dim j < \dim i$ for every non-identity morphism $\delta : j \rightarrow i$.

Definition

A direct category \mathcal{I} is a small category equipped with a dimension function $\dim : \text{ob } \mathcal{I} \rightarrow \text{Ord}$ to the class of ordinals such that $\dim j < \dim i$ for every non-identity morphism $\delta : j \rightarrow i$.

Example

- Any **discrete** category S .

Direct Categories

Definition

A direct category \mathcal{I} is a small category equipped with a dimension function $\dim : \text{ob } \mathcal{I} \rightarrow \text{Ord}$ to the class of ordinals such that $\dim j < \dim i$ for every non-identity morphism $\delta : j \rightarrow i$.

Example

- Any discrete category S .
- The category \mathbb{G} of globes

$$[0] \begin{array}{c} \xrightarrow{s} \\ \xleftarrow{t} \end{array} [1] \begin{array}{c} \xrightarrow{s} \\ \xleftarrow{t} \end{array} [2] \begin{array}{c} \xrightarrow{s} \\ \xleftarrow{t} \end{array} \dots$$

$$ss = ts$$

$$st = tt$$

Definition

A direct category \mathcal{I} is a small category equipped with a dimension function $\dim : \text{ob } \mathcal{I} \rightarrow \text{Ord}$ to the class of ordinals such that $\dim j < \dim i$ for every non-identity morphism $\delta : j \rightarrow i$.

Example

- Any discrete category S .
- The category \mathbb{G} of globes
- The category Δ_{inj} of **simplices** and face maps.

$$[0] \begin{array}{c} \xrightarrow{\delta_0} \\ \xleftarrow{\delta_1} \end{array} [1] \begin{array}{c} \xrightarrow{\delta_0} \\ \xleftarrow{\delta_2} \end{array} [2] \begin{array}{c} \xrightarrow{\delta_0} \\ \xleftarrow{\delta_3} \end{array} \cdots \quad \delta_i \delta_j = \delta_{j+1} \delta_i \quad (i \leq j)$$

Direct Categories

Definition

A direct category \mathcal{I} is a small category equipped with a dimension function $\dim : \text{ob } \mathcal{I} \rightarrow \text{Ord}$ to the class of ordinals such that $\dim j < \dim i$ for every non-identity morphism $\delta : j \rightarrow i$.

Example

- Any discrete category S .
- The category \mathbb{G} of globes
- The category Δ_{inj} of simplices and face maps.

Notation

We denote by \mathcal{I}_α the full subcategory on objects of dimension at most α with the obvious dimension function.

The inductive data

We define by transfinite recursion on $\alpha \leq \sup\{\dim i : i \in \mathcal{I}\}$

- a class $\text{Sig}_\alpha(\mathcal{I})$ of (generalised) signatures of dimension α ,

The inductive data

We define by transfinite recursion on $\alpha \leq \sup\{\dim i : i \in \mathcal{I}\}$

- a class $\text{Sig}_\alpha(\mathcal{I})$ of (generalised) signatures of dimension α ,
- restriction functions $\text{Sig}_\alpha(\mathcal{I}) \xrightarrow{(-)_\beta} \text{Sig}_\beta(\mathcal{I})$ for $\beta < \alpha$

The inductive data

We define by transfinite recursion on $\alpha \leq \sup\{\dim i : i \in \mathcal{I}\}$

- a class $\text{Sig}_\alpha(\mathcal{I})$ of (generalised) signatures of dimension α ,
- restriction functions $\text{Sig}_\alpha(\mathcal{I}) \xrightarrow{(-)_\beta} \text{Sig}_\beta(\mathcal{I})$ for $\beta < \alpha$
- an adjunction for every signature Σ of dimension α

$$[\mathcal{I}_\alpha^{\text{op}}, \text{Set}] \begin{array}{c} \xrightarrow{\text{Cptd}_\Sigma} \\ \perp \\ \xleftarrow{\text{Term}_\Sigma} \end{array} \text{Comp}_\Sigma$$

The inductive data

We define by transfinite recursion on $\alpha \leq \sup\{\dim i : i \in \mathcal{I}\}$

- a class $\text{Sig}_\alpha(\mathcal{I})$ of (generalised) signatures of dimension α ,
- restriction functions $\text{Sig}_\alpha(\mathcal{I}) \xrightarrow{(-)_\beta} \text{Sig}_\beta(\mathcal{I})$ for $\beta < \alpha$
- an adjunction for every signature Σ of dimension α

$$[\mathcal{I}_\alpha^{\text{op}}, \text{Set}] \begin{array}{c} \xrightarrow{\text{Cptd}_\Sigma} \\ \perp \\ \xleftarrow{\text{Term}_\Sigma} \end{array} \text{Comp}_\Sigma$$

- truncation functors $\text{Comp}_\Sigma \xrightarrow{\text{tr}_\beta^\Sigma} \text{Comp}_{\Sigma_\beta}$ for every $\beta < \alpha$

Definition

A signature Σ of dimension α consists of

- signatures Σ_β for $\beta < \alpha$ such that $(\Sigma_\beta)_\gamma = \Sigma_\gamma$,

Definition

A signature Σ of dimension α consists of

- signatures Σ_β for $\beta < \alpha$ such that $(\Sigma_\beta)_\gamma = \Sigma_\gamma$,
- sets Σ_i of function symbols for $\dim i = \alpha$,

Definition

A signature Σ of dimension α consists of

- signatures Σ_β for $\beta < \alpha$ such that $(\Sigma_\beta)_\gamma = \Sigma_\gamma$,
- sets Σ_i of function symbols for $\dim i = \alpha$,
- for every function symbol $f \in \Sigma_i$,

Definition

A signature Σ of dimension α consists of

- signatures Σ_β for $\beta < \alpha$ such that $(\Sigma_\beta)_\gamma = \Sigma_\gamma$,
- sets Σ_i of function symbols for $\dim i = \alpha$,
- for every function symbol $f \in \Sigma_i$,
 - an arity $B_f \in [\mathcal{I}^{\text{op}}, \text{Set}]$

Definition

A signature Σ of dimension α consists of

- signatures Σ_β for $\beta < \alpha$ such that $(\Sigma_\beta)_\gamma = \Sigma_\gamma$,
- sets Σ_i of function symbols for $\dim i = \alpha$,
- for every function symbol $f \in \Sigma_i$,
 - an arity $B_f \in [\mathcal{I}^{\text{op}}, \text{Set}]$
 - for non-identity $\delta : j \rightarrow i$, a boundary term

$$\delta^* f \in \text{Term}_{\Sigma_{\dim j}, j} \text{Cptd}_{\Sigma_{\dim j}}(\text{tr}_{\dim j} B_f)$$

satisfying that $(\delta')^*(\delta^* f) = (\delta\delta')^* f$

The restriction functions are the obvious projections.

Definition

A Σ -computad C consists of

- Σ_β -computads C_β for $\beta < \alpha$ such that $\text{tr}_\gamma^{\Sigma_\beta} C_\beta = C_\gamma$,

Definition

A Σ -computad C consists of

- Σ_β -computads C_β for $\beta < \alpha$ such that $\text{tr}_\gamma^{\Sigma_\beta} C_\beta = C_\gamma$,
- a set V_i^C of generators for $\dim i = \alpha$,

Definition

A Σ -computad C consists of

- Σ_β -computads C_β for $\beta < \alpha$ such that $\text{tr}_\gamma^{\Sigma_\beta} C_\beta = C_\gamma$,
- a set V_i^C of generators for $\dim i = \alpha$,
- gluing functions $V_i^C \rightarrow \text{Term}_{\Sigma_{\dim j, j}}(C_{\dim j})$ for non-identity $\delta : j \rightarrow i$ such that

$$(\delta')^* \phi_\delta^C = \phi_{\delta\delta'}^C$$

Definition

A Σ -computad C consists of

- Σ_β -computads C_β for $\beta < \alpha$ such that $\text{tr}_\gamma^{\Sigma_\beta} C_\beta = C_\gamma$,
- a set V_i^C of generators for $\dim i = \alpha$,
- gluing functions $V_i^C \rightarrow \text{Term}_{\Sigma_{\dim j}, j}(C_{\dim j})$ for non-identity $\delta : j \rightarrow i$ such that

$$(\delta')^* \phi_\delta^C = \phi_{\delta\delta'}^C$$

Each presheaf $X : \mathcal{I}_\alpha^{\text{op}} \rightarrow \text{Set}$ defines a computad by

$$(\text{Cptd}_\Sigma X)_\beta = \text{Cptd}_{\Sigma_\beta} \text{tr}_\beta X$$

$$V_i^{\text{Cptd}_\Sigma X} = X_i$$

$$\phi_\delta^{\text{Cptd}_\Sigma X} = \eta_{\Sigma_{\dim j}} \circ \delta^*$$

Definition

The lower dimensional terms of a computad C are those of the computads C_β . The set $\text{Term}_{\Sigma,i} C$ for $\dim i = \alpha$ is inductively generated by

- $\text{var} : V_i^C \rightarrow \text{Term}_{\Sigma,i} C$
- $\text{comp} : \sum_{f \in \Sigma_i} \text{Comp}_\Sigma(\text{Cptd}_\Sigma B_f, C) \rightarrow \text{Term}_{\Sigma,i} C$

Terms form a presheaf by letting for $\delta : j \rightarrow i$

$$\begin{aligned}\delta^*(\text{var } v) &= \phi_\delta^C(v) \\ \delta^*(\text{comp}[f, \tau]) &= \text{Term}_{\Sigma_{\dim j, j}}(\tau_{\dim j})(\delta^* f)\end{aligned}$$

Definition

A morphism $\sigma : D \rightarrow C$ consists of

- morphisms $\sigma_\beta : D_\beta \rightarrow C_\beta$ for $\beta < \alpha$ such that $\text{tr}_{\gamma}^{\Sigma_\beta} \sigma_\beta = \sigma_\gamma$,
- functions $\sigma_i : V_i^D \rightarrow \text{Term}_{\Sigma, i}(C)$ for $\dim i = \alpha$ satisfying for $\delta : j \rightarrow i$ that

$$\begin{array}{ccc} V_i^D & \xrightarrow{\sigma_i} & \text{Term}_{\Sigma, i} C \\ \downarrow \phi_\delta^D & & \downarrow \delta^* \\ \text{Term}_{\Sigma_{\dim j}, j} D & \xrightarrow{\sigma_{\dim j}} & \text{Term}_{\Sigma_{\dim j}, j} C \end{array}$$

Definition

A morphism $\sigma : D \rightarrow C$ consists of

- morphisms $\sigma_\beta : D_\beta \rightarrow C_\beta$ for $\beta < \alpha$ such that $\text{tr}_{\gamma}^{\Sigma_\beta} \sigma_\beta = \sigma_\gamma$,
- functions $\sigma_i : V_i^D \rightarrow \text{Term}_{\Sigma, i}(C)$ for $\dim i = \alpha$ satisfying for $\delta : j \rightarrow i$ that

$$\begin{array}{ccc} V_i^D & \xrightarrow{\sigma_i} & \text{Term}_{\Sigma, i} C \\ \downarrow \phi_\delta^D & & \downarrow \delta^* \\ \text{Term}_{\Sigma_{\dim j}, j} D & \xrightarrow{\sigma_{\dim j}} & \text{Term}_{\Sigma_{\dim j}, j} C \end{array}$$

We will say that $\sigma : D \rightarrow C$ is **variable-to-variable** when σ_β are variable-to-variable and σ_i factors via var .

Given a morphism $\sigma : C \rightarrow D$, we define composition and the action on terms recursively by

$$\sigma \circ \tau = (\sigma_\beta \circ \tau_\beta, \text{Term}_{\Sigma,i}(\sigma) \circ \tau_i)$$

where

$$\begin{aligned}\text{Term}_{\Sigma,i}(\sigma)(\text{var } v) &= \sigma_i(v) \\ \text{Term}_{\Sigma,i}(\sigma)(\text{comp}[f, \tau]) &= \text{comp}[f, \sigma \circ \tau]\end{aligned}$$

The term adjunction

- The unit includes generators into terms

$$\begin{aligned}\eta_{\Sigma, X} : X &\rightarrow \text{Term}_{\Sigma} \text{Cptd}_{\Sigma} X \\ x &\mapsto \text{var } x\end{aligned}$$

The term adjunction

- The unit includes generators into terms

$$\eta_{\Sigma, X} : X \rightarrow \text{Term}_{\Sigma} \text{Cptd}_{\Sigma} X$$
$$x \mapsto \text{var } x$$

- The counit identifies terms with generators

$$\varepsilon_{\Sigma, C} : \text{Cptd}_{\Sigma} \text{Term}_{\Sigma} C \rightarrow C$$
$$\varepsilon_{\Sigma, C, \beta} = \epsilon_{\Sigma_{\beta}, C_{\beta}}$$
$$\varepsilon_{\Sigma, C, i} : V_i^{\text{Cptd}_{\Sigma} \text{Term}_{\Sigma} C} = \text{Term}_{\Sigma} C$$

Definition

The term monad $(M_\Sigma, \eta_\Sigma, \mu_\Sigma)$ is the monad on $[\mathcal{I}_\alpha^{\text{op}}, \text{Set}]$ induced by the term adjunction. The category Alg_Σ is the category of M_Σ -algebras.

Definition

The term monad $(M_\Sigma, \eta_\Sigma, \mu_\Sigma)$ is the monad on $[\mathcal{I}_\alpha^{\text{op}}, \text{Set}]$ induced by the term adjunction. The category Alg_Σ is the category of M_Σ -algebras.

Algebras $\mathbb{X} = (X, u^\mathbb{X})$ are pairs of a presheaf and a morphism

$$u^\mathbb{X} : \text{Term}_\Sigma \text{Cptd}_\Sigma X \rightarrow X$$

satisfying two axioms.

Definition

The term monad $(M_\Sigma, \eta_\Sigma, \mu_\Sigma)$ is the monad on $[\mathcal{I}_\alpha^{\text{op}}, \text{Set}]$ induced by the term adjunction. The category Alg_Σ is the category of M_Σ -algebras.

Algebras $\mathbb{X} = (X, u^\mathbb{X})$ are pairs of a presheaf and a morphism

$$u^\mathbb{X} : \text{Term}_\Sigma \text{Cptd}_\Sigma X \rightarrow X$$

satisfying two axioms. They give rise for every $f \in \Sigma_i$ to functions

$$f^\mathbb{X} : [\mathcal{I}_\alpha^{\text{op}}, \text{Set}](B_f, X) \rightarrow X;$$
$$f^\mathbb{X}(\tau) = u^\mathbb{X}(\text{comp}[f, \text{Cptd}_\Sigma \tau])$$

and they are determined by them.

Table of Contents

1 Motivation

2 Definitions


3 Examples

4 Results

Notation

We will let $\text{ssSet} = [\Delta_{\text{inj}}^{\text{op}}, \text{Set}]$ and


- $\Delta[n]$ the representable semisimplicial set



Notation

We will let $\text{ssSet} = [\Delta_{\text{inj}}^{\text{op}}, \text{Set}]$ and


- $\Delta[n]$ the representable semisimplicial set
- $\partial\Delta[n]$ its boundary



Notation

We will let $\text{ssSet} = [\Delta_{\text{inj}}^{\text{op}}, \text{Set}]$ and


- $\Delta[n]$ the representable semisimplicial set
- $\partial\Delta[n]$ its boundary
- $\Lambda^k[n]$ its k -th horn



Notation

We will let $\text{ssSet} = [\Delta_{\text{inj}}^{\text{op}}, \text{Set}]$ and

- $\Delta[n]$ the representable semisimplicial set
- $\partial\Delta[n]$ its boundary
- $\Lambda^k[n]$ its k -th horn



Definition

An algebraic semisimplicial Kan complex \mathbb{X} is a semisimplicial set X together with for $n > 0$ and $0 \leq k \leq n$ a choice of lift for


$$\begin{array}{ccc} \Lambda^k[n] & \xrightarrow{\sigma} & X \\ \downarrow & \nearrow \text{dashed} & \\ \Delta[n] & & \end{array}$$

Kan complexes

Notation

We will let $\text{ssSet} = [\Delta_{\text{inj}}^{\text{op}}, \text{Set}]$ and

- $\Delta[n]$ the representable semisimplicial set
- $\partial\Delta[n]$ its boundary
- $\Lambda^k[n]$ its k -th horn



Definition

An algebraic semisimplicial Kan complex \mathbb{X} is a semisimplicial set X together with for $n > 0$ and $0 \leq k \leq n$ a choice of lift for

$$\begin{array}{ccc} \Lambda^k[n] & \xrightarrow{\sigma} & X \\ \downarrow & \nearrow \text{dashed} & \\ \Delta[n] & & \end{array}$$

$$\text{face}_{k,n} : \text{ssSet}(\Lambda^k[n], X) \rightarrow X_{n-1}$$

$$\text{fill}_{k,n} : \text{ssSet}(\Lambda^k[n], X) \rightarrow X_n$$

Kan complexes are algebras of a signature defined by

$$\Sigma_{\text{Kan},[n]} = \{\text{face}_{n+1,k} : 0 \leq k \leq n+1\} \cup \{\text{fill}_{k,n} : n > 0, 0 \leq k \leq n\}$$

$$B_{\text{face}_{k,n+1}} = \Lambda^k[n+1]$$

$$\delta^* \text{face}_{k,n+1} = \text{var}(\delta_k \delta)$$

$$B_{\text{fill}_{k,n}} = \Lambda^k[n]$$

$$\delta^* \text{fill}_{k,n} = \begin{cases} \text{face}_{k,n}, & \text{if } \delta = \delta_k \\ \text{var } \delta, & \text{otherwise.} \end{cases}$$

Kan Complexes

Kan complexes are algebras of a signature defined by

$$\Sigma_{\text{Kan},[n]} = \{\text{face}_{n+1,k} : 0 \leq k \leq n+1\} \cup \{\text{fill}_{k,n} : n > 0, 0 \leq k \leq n\}$$

$$B_{\text{face}_{k,n+1}} = \Lambda^k[n+1]$$

$$\delta^* \text{face}_{k,n+1} = \text{var}(\delta_k \delta)$$

$$B_{\text{fill}_{k,n}} = \Lambda^k[n]$$

$$\delta^* \text{fill}_{k,n} = \begin{cases} \text{face}_{k,n}, & \text{if } \delta = \delta_k \\ \text{var } \delta, & \text{otherwise.} \end{cases}$$

Remark

Algebraic semisimplicial Kan complexes carry a model structure equivalent to spaces⁵. We will discuss its cofibrations shortly.

⁵Bourke and Henry, *Algebraically cofibrant and fibrant object revisited*

Globular Higher Categories

Definition

We let $\text{Glob} = [\mathbb{G}^{\text{op}}, \text{Set}]$ the category of globular sets. We let Bat be the family of globular sets representing the strict ω -category monad.



Definition

We let $\text{Glob} = [\mathbb{G}^{\text{op}}, \text{Set}]$ the category of globular sets. We let Bat the family of globular sets representing the strict ω -category monad.

The signature $\Sigma_{\omega \text{ cat}}$ for weak ω -categories⁶

- has no operations of dimension 0,
- has as operations of dimension $n + 1$ triples (B, a, b) where
 - $B \in \text{Bat}$ of dimension at most $n + 1$,
 - $a, b \in \text{Term}_n(B)$ that “cover” the n -dimensional source and target of B respectively, and have common source and target.



⁶Dean et al., *Computads for weak ω -categories as an inductive type*

Table of Contents

1 Motivation

2 Definitions

3 Examples

4 Results

Lemma

The subcategory $\text{Comp}_{\Sigma}^{\text{var}}$ of variable-to-variable morphisms has a terminal computad $\mathbb{1}_{\Sigma}$.

Generators of the terminal computad represent the “shapes” of generators in the sense that

$$V_i^C \cong \coprod_{p \in V_i^{\mathbb{1}_{\Sigma}}} \text{Comp}_{\Sigma}^{\text{var}}(|p|, C).$$

Lemma

The subcategory $\text{Comp}_{\Sigma}^{\text{var}}$ of variable-to-variable morphisms has a terminal computad $\mathbb{1}_{\Sigma}$.

Generators of the terminal computad represent the “shapes” of generators in the sense that


$$V_i^C \cong \coprod_{p \in V_i^{\mathbb{1}_{\Sigma}}} \text{Comp}_{\Sigma}^{\text{var}}(|p|, C).$$


Theorem

Let Plex_{Σ} the full subcategory of $\text{Comp}_{\Sigma}^{\text{var}}$ on the computads $|p|$. The subcategory inclusion induces an equivalence


$$\text{Comp}_{\Sigma}^{\text{var}} \cong [\text{Plex}_{\Sigma}^{\text{op}}, \text{Set}]$$

Free Algebras






- K_{Σ} is fully faithful




- K_{Σ} is fully faithful
- $\text{Cptd}_{\Sigma}^{\text{var}}$ is fully faithful



- K_{Σ} is fully faithful
- $\text{Cptd}_{\Sigma}^{\text{var}}$ is fully faithful
- Morphisms $\text{Fr}_{\Sigma} C \rightarrow \mathbb{X}$ amount to functions

$$V_i^C \rightarrow X_i$$

compatible with the
gluing functions.



- K_{Σ} is fully faithful
- $\text{Cptd}_{\Sigma}^{\text{var}}$ is fully faithful
- Morphisms $\text{Fr}_{\Sigma} C \rightarrow \mathbb{X}$ amount to functions

$$V_i^C \rightarrow X_i$$

compatible with the
gluing functions.

Computads as cofibrant algebras

Let $\mathbb{D}^i = \text{Fr}_\Sigma \mathcal{I}(-, i)$ and $\partial\mathbb{D}^i$ its boundary. The inclusions $\partial\mathbb{D}_\Sigma^i \subseteq \mathbb{D}_\Sigma^i$ cofibrantly generate a weak factorisation system on the category of algebras.

Remark

This is the (cofibration, trivial fibration) wfs for Kan complexes and for strict ω -categories.

Computads as cofibrant algebras

Let $\mathbb{D}^i = \text{Fr}_\Sigma \mathcal{I}(-, i)$ and $\partial \mathbb{D}^i$ its boundary. The inclusions $\partial \mathbb{D}_\Sigma^i \subseteq \mathbb{D}_\Sigma^i$ cofibrantly generate a weak factorisation system on the category of algebras.

Theorem

Free algebras on a computad are cofibrant.

Remark

This is the (cofibration, trivial fibration) wfs for Kan complexes and for strict ω -categories.

Computads as cofibrant algebras

Let $\mathbb{D}^i = \text{Fr}_\Sigma \mathcal{I}(-, i)$ and $\partial \mathbb{D}^i$ its boundary. The inclusions $\partial \mathbb{D}_\Sigma^i \subseteq \mathbb{D}_\Sigma^i$ cofibrantly generate a weak factorisation system on the category of algebras.

Theorem

Free algebras on a computad are cofibrant.

Theorem

The comonad $\text{Fr}_\Sigma \text{Und}_\Sigma : \text{Alg}_\Sigma \rightarrow \text{Alg}_\Sigma$ is the universal cofibrant replacement⁷ of the wfs.

Remark

This is the (cofibration, trivial fibration) wfs for Kan complexes and for strict ω -categories.

⁷Garner, “Homomorphisms of higher categories”

Computads as cofibrant algebras

Let $\mathbb{D}^i = \text{Fr}_\Sigma \mathcal{I}(-, i)$ and $\partial \mathbb{D}^i$ its boundary. The inclusions $\partial \mathbb{D}_\Sigma^i \subseteq \mathbb{D}_\Sigma^i$ cofibrantly generate a weak factorisation system on the category of algebras.

Theorem

Free algebras on a computad are cofibrant, and vice versa.







Theorem

The comonad $\text{Fr}_\Sigma \text{Und}_\Sigma : \text{Alg}_\Sigma \rightarrow \text{Alg}_\Sigma$ is the universal cofibrant replacement⁷ of the wfs.

Remark

This is the (cofibration, trivial fibration) wfs for Kan complexes and for strict ω -categories.

⁷Garner, “Homomorphisms of higher categories”

-  Batanin, M. A. “Computads for finitary monads on globular sets”. In: *Higher Category Theory*. Vol. 230. Contemporary Mathematics. American Mathematical Society, 1998.
-  Bourke, J. and R. Garner. “Monads and theories”. In: *Advances in Mathematics* 351 (2019). arXiv: 1805.04346.
-  Bourke, J. and S. Henry. *Algebraically cofibrant and fibrant object revisited*. 2020. arXiv: 2005.05384.
-  Dean, C. J., E. Finster, I. Markakis, D. Reutter, and J. Vicary. *Computads for weak ω -categories as an inductive type*. 2022. arXiv: 2208.08719.
-  Garner, R. “Homomorphisms of higher categories”. In: *Advances in Mathematics* 224.6 (2010). arXiv: 0810.4450.
-  Street, R. “Limits indexed by category-valued 2-functors”. In: *Journal of Pure and Applied Algebra* 8.2 (1976).